
Assessing the Effects of Altered Marine Nitrogen Flow on Ecosystem Health and Biodiversity Across Various Ecoregions

Yourfee Kennedy, Biosphere Reserves Management, MSc. Eberswalde University for Sustainable Development, Germany

Abstract

The study delves into how changes in marine nitrogen flows affect the health and variety of life within different environmental regions, employing Geographic Information System (GIS) techniques for analysis. It focuses on mapping out the distribution, categorizing the densities, measuring the intensity, and identifying the geographical areas affected by marine nitrogen changes. Results indicate a varied pattern in how nitrogen moves through different marine environments, pointing out how crucial spatial variations in nitrogen accumulation are for the health of ecosystems and the conservation of biodiversity. This research places a spotlight on the increasing influence of human activities on the balance of nutrients in the ocean, particularly in areas where there has been a significant rise in the levels of nitrogen flow. It stresses the urgent need for creating specific conservation plans aimed at reducing the negative impacts of nutrient pollution on marine life and ensuring the preservation of diversity in these habitats.

Keywords: Marine ecosystems, Nitrogen flux, Biodiversity, Geographic Information System (GIS), Ecoregions, Spatial analysis, Conservation, Human activities, Nutrient pollution, Eutrophication, Sustainable management, and Ecological modeling.

1 Introduction

Marine ecosystems play a crucial role in upholding global biodiversity and facilitating essential ecological processes. Among the multitude of factors impacting the health of marine ecosystems, nitrogen flux emerges as a key driver that influences primary productivity, nutrient cycling, and biodiversity (Nilles and Conley 2001, Lehmann et al. 2005, Conley et al., 2009). Nitrogen, being a vital nutrient, is intricately associated with various biological mechanisms within marine settings. Nonetheless, modifications in the flow of nitrogen in marine environments, triggered by human activities and changes in the environment, present substantial obstacles to ecosystem well-being and the preservation of biodiversity (Gruber & Galloway, 2008, Galloway et al. 2003). Comprehending the consequences of these alterations across different ecoregions is crucial for the efficient management of the environment and conservation endeavours. This research endeavor is focused on investigating the impacts of modified marine nitrogen flux on ecosystem well-being and biodiversity in diverse ecoregions. Through the application of Geographic Information System (GIS) methodologies, the spatial distribution of nitrogen flux in marine settings is examined, along with an evaluation of its consequences on ecosystem dynamics and biodiversity. The primary objective of this study is to shed light on the diverse reactions of marine ecosystems to altered nitrogen inputs by analysing nitrogen flux dynamics across various ecoregions. In doing so, valuable insights into the factors driving ecological transformations are provided, offering guidance for the development of conservation strategies aimed at fostering sustainable marine management (Fenn et al. 2010, Collins et al. 2011, Galloway et al. 2004).

2 Literature Review

Past studies have played a pivotal role in demonstrating how nitrogen is a key factor in the development and diversity of marine ecosystems, as documented by Conley and his colleagues (IPCC 2007; Conley et al. 2009). These research efforts have shed light on nitrogen's crucial influence over primary productivity and the sustenance of varied marine life. The changes in nitrogen levels, whether from natural events or anthropogenic sources, are known to significantly impact the ecological architecture and its operations (Howarth & Marino, 2006; Gruber & Galloway, 2008). For instance, an overabundance of nitrogen stemming from farm runoff, urban waste, and air pollution has been identified as a cause of eutrophication, detrimental algae growth, and a decrease in oxygen levels in oceanic habitats (Diaz & Rosenberg, 2008; Paerl & Otten, 2013; Clark and Tilman 2008; Fernandez et al. 2010; Driscoll et al. 2012; Groffman et al. 2012). While the effects of nitrogen on coastal landscapes have been thoroughly examined, the exploration into how nitrogen dynamics influence various marine ecosystems worldwide remains limited. It's critical to comprehend the spatial and temporal variations of nitrogen within multiple aquatic environments to foretell how ecosystems reacts to changes in their environment and to craft precise conservation strategies (Rabalais et al., 2009; Seitzinger et al., 2010). The use of Geographic Information System (GIS) technology stands out as an influential means for assessing nitrogen distribution and biodiversity data. This allows investigators to delve into the intricate interactions between environmental factors and biological distributions (Lehner & Döll, 2004; Lillesand & Kiefer, 2000; Tierney et al. 2001; Christopher et al. 2008; Shibata et al. 2013; Piatek et al. 2005).

3 Methodology

Utilizing Geographic Information System (GIS) methods, this study investigates how changes in marine nitrogen transportation affect ecosystem well-being and biodiversity across varied ecological regions. The research is centered on applying QGIS software to carry out spatial examinations of marine nitrogen movement through polygon shapefiles within distinct ecological areas. The process involves gathering polygon shapefile data from reputable sources, followed by their examination and analysis to measure the patterns of nitrogen movement and to evaluate their spatial arrangement throughout the area under study. The approach encompasses multiple phases, like data preprocessing, conducting spatial analysis, and interpreting the findings. At the outset, polygon shapefile data gets imported into the QGIS setting, where it undergoes preprocessing steps that involve cleaning the data, adjusting projections, and normalizing attributes. Following this, techniques for spatial analysis are employed to investigate the spatial distribution patterns of marine nitrogen flux, which include analysis of hotspots, studying spatial autocorrelation, and employing interpolation techniques (referenced by Hoekstra and Molnar, 2010, Bivand et al., 2013; Olaya, 2004). Analyzing spatial data unveils the unequal distribution of nitrogen flux within various ecoregions, pinpointing zones with elevated and diminished nitrogen concentrations. Overlaying data concerning nitrogen flux with biodiversity indices or ecological metrics serves the purpose of shedding light on how nitrogen movements influence the health of ecosystems. Implementations of statistical analyses, like correlation analyses and regression models, are anticipated to measure the relationship between parameters of nitrogen flux and indicators of biodiversity (cited by Fredston-Hermann et al., 2016, Fielding & Bell, 1997; Getis & Ord, 1992).

4 Results

4.1 Spatial Distribution of Marine Nitrogen Flow by Ecoregion




Figure 1. Shows the distribution of Marine Nitrogen flow by Ecoregions.

4.2 Population Density Categorization Across Ecoregions




Figure 2. Shows the density categorization of marine Nitrogen Flow across Ecoregions.

4.3 Nitrogen Flow Intensity Across Ecoregions




Figure 3. Shows the intensity of Marine Nitrogen Flow across Ecoregions.

4.4 Marine Nitrogen Flow by Geographic Realms




Figure 4. Shows the realms of Marine Nitrogen Flow.

5 Discussion

Figure 1's illustration of the geographical distribution of nitrogen circulation within various marine ecoregions uncovers uneven patterns spanning across different aquatic environments. Regions like the North Sea, Line Islands, and Levantine Sea display differing nitrogen circulation levels, mirroring the regional variances in nutrient dynamics and the factors driving these environments. These observations are consistent with prior research that highlighted the critical role of spatial differences in nitrogen deposits for the functioning of ecosystems and the preservation of biodiversity (Worm and Lotze, 2016; Conley et al., 2009; Rabalais et al., 2009). Pinpointing ecoregions where nitrogen circulation is high is crucial for directing conservation actions and developing management plans aimed to lessen nutrient pollution's effects on marine ecosystems. The mapping out of nitrogen circulation intensity across ecoregions, as shown in Figure 2, points out the time-based trends in nitrogen deposits and their repercussions on the wellbeing of ecosystems. The yearly escalation of nitrogen circulation intensity across all regions emphasizes the escalating impact of human-related activities on marine nutrient dynamics (Howarth & Marino, 2006). Regions like the Adriatic and Aegean Seas are witnessing noticeable upticks in nitrogen circulation intensity, pointing to potential ecological dangers stemming from nutrient overload and eutrophication (Drupp et al., 2011; Diaz & Rosenberg, 2008). These discoveries point to the pressing necessity for sustainable management approaches to confront nutrient pollution and protect marine biodiversity in at-risk ecoregions (Urban et al., 2016).

6 Conclusion

This research utilizes Geographic Information System (GIS) technology to explore how changes in marine nitrogen flow affects the health and biodiversity of ecosystems within different ecoregions. By conducting a spatial analysis on the flow patterns of nitrogen, the study sheds light on its spatial arrangement, categorization by density, the intensity of flow, and the areas affected by marine nitrogen movement. This offers valuable insights into the problematic nature of nutrient pollution for ecological systems. The outcomes emphasize the critical nature of spatial differences in nitrogen deposits on the functioning of ecosystems and underline the necessity for specially tailored conservation plans to reduce the detrimental consequences of nutrient pollution in marine environments. This research merges GIS-based spatial analysis with ecological modeling techniques, enhancing our comprehension of how nitrogen dynamics interact with the health of ecosystems. Such understanding is crucial for directing conservation activities aimed towards the sustainable management of marine environments and the preservation of their biodiversity (Cavanaugh et al., 2014).

7 References

- Aber JD, Ollinger SV, Driscoll CT, Likens GE, Holmes RT, Freuder RJ, Goodale CL. Inorganic nitrogen losses from a forested ecosystem in response to physical, chemical, biotic, and climatic perturbations. *Ecosystems*. 2002; 5:648–658. [Google Scholar]
- Anderson CB, Celis-Diez JL, Bond BJ, Pastur GM, Little C, Armesto JJ, Ghera C, Austin A, et al. Progress in creating a joint research agenda that allows networked long-term socio-ecological research in southern South America: Addressing crucial technological and human capacity gaps limiting its application in Chile and Argentina. *Austral Ecology*. 2012;37:529–536. [Google Scholar]
- Beier C, Emmett B, Gundersen P, Tietema A, Peñuelas J, Estiarte M, Gordon C, Gorissen A, et al. Novel approaches to study climate change effects on terrestrial ecosystems in the field: Drought and passive nighttime warming. *Ecosystems*. 2004;7:583–597. [Google Scholar]
- Bergström A, Jansson M. Atmospheric nitrogen deposition has caused nitrogen enrichment and eutrophication of lakes in the northern hemisphere. *Global Change Biology*. 2006;12:635–643. [Google Scholar]
- Billen G, Thieu V, Garnier J, Silvestre M. Modelling the N cascade in regional watersheds: The case study of the Seine, Somme and Scheldt rivers. *Agriculture, Ecosystems and Environments*. 2009;133:234–246. [Google Scholar]
- Blenckner T, Adrian R, Livingstone DM, Jennings E, Weyhenmeyer GA, George DG, Jankowski T, Järvinen M, et al. Large-scale climatic signatures in lakes across Europe: A meta-analysis. *Global Change Biology*. 2007; 13:1314–1326. [Google Scholar]
- Bobbink, R., and J.P. Hettelingh. 2011. Review and revision of empirical critical loads and dose-response relationships, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM).
- Bobbink R, Hicks K, Galloway J, Spranger T, Alkemade R, Ashmore M, Bustamante M, Cinderby S, et al. Global assessment of nitrogen deposition effects on terrestrial plant diversity: A synthesis. *Ecological Applications*. 2010; 20:30–59. [PubMed] [Google Scholar]
- Boberg JB, Finlay RD, Stenlid J, Lindahl BD. Fungal C translocation restricts N-mineralization in heterogeneous environments. *Functional Ecology*. 2010; 24:454–459. [Google Scholar]
- Bowman WD, Cleveland CC, Halada L, Hresko LJ, Baron JS. Negative impact of nitrogen deposition on soil buffering capacity. *Nature Geoscience*. 2008; 1:767–770. [Google Scholar]
- Boyer EW, Howarth RW. The nitrogen cycle at regional to global scales. Dordrecht: Kluwer; 2002. [Google Scholar]

Camarero L, Catalan J. Atmospheric phosphorus deposition may cause lakes to revert from phosphorus limitation back to nitrogen limitation. *Nature Communications*. 2012;3:1118. [PubMed] [Google Scholar]

Campbell JL, Mitchell MJ, Groffman PM, Christenson LM. Winter in northeastern North America: An often overlooked but critical period for ecological processes. *Frontiers in Ecology and the Environment*. 2005;3:314–322. [Google Scholar]

Cape JN, Eerden LJVD, Sheppard LJ, Leith ID, Sutton MA. Evidence for changing the critical level for ammonia. *Environmental Pollution*. 2009;157:1033–1037. [PubMed] [Google Scholar]

Carpenter SR, Cole JJ, Hodgson JR, Kitchell JF, Pace ML, Bade D, Cottingham KL, Essington TE, Houder JN, et al. Trophic cascades, nutrients, and lake productivity: Whole-lake experiments. *Ecological Monographs*. 2001;71:163–186. [Google Scholar]

Casson NJ, Eimers MC, Buttle JM. The Contribution of Rain-On-Snow Events to Nitrate Export in the Forested Landscape of South-Central Ontario, Canada. *Hydrological Processes*. 2010;24:1985–1993. [Google Scholar]

Chestnut TJ, McDowell WH. C and N dynamics in the riparian and hyporheic zones of a tropical stream, Luquillo mountains, Puerto Rico. *Journal of the North American Benthological Society*. 2000;19:199–214. [Google Scholar]

Christopher SF, Shibata H, Ozawa M, Nakagawa Y, Mitchell MJ. The effect of soil freezing on N cycling: Comparison of two headwater subcatchments with different vegetation and snowpack conditions in the northern Hokkaido Island of Japan. *Biogeochemistry*. 2008;88:15–30. [Google Scholar]

Clark CM, Tilman D. Loss of plant species after chronic low-level nitrogen deposition to prairie grasslands. *Nature*. 2008;451:712–715. [PubMed] [Google Scholar]

Collins SL, Carpenter SR, Swinton SM, Orenstein DE, Childers DL, Gragson TL, Grimm NB, Grove JM, et al. An integrated conceptual framework for long-term social–ecological research. *Frontiers in Ecology and the Environment*. 2011;9:351–357. [Google Scholar]

Compton JE, Harrison JA, Dennis RL, Greaver TL, Hill BH, Jordan SJ, Walker H, Campbell HV. Ecosystem services altered by human changes in the nitrogen cycle: A new perspective for US decision making. *Ecological Letters*. 2011;14:804–815. [PubMed] [Google Scholar]

Curtis CJ, Heaton TEH, Simpson GL, Evans CD, Shilland J, Turner S. Dominance of biologically produced nitrate in upland waters of Great Britain indicated by stable isotopes. *Biogeochemistry*. 2012;111:535–554. [Google Scholar]

Curtis, C.J., R.W. Battarbee, D.T. Monteith, and E.M. Shilland. 2014. The future of upland water ecosystems of the UK in the 21st century: A synthesis. *Ecological Indicators* 37(Part B): 412–430.

- Cusack DF, Silver WL, Torn MS, McDowell WH. Effects of nitrogen additions on above- and belowground carbon dynamics in two tropical forests. *Biogeochemistry*. 2011;104:203–225. [Google Scholar]
- Davidson EA. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nature Geoscience*. 2009;2:659–662. [Google Scholar]
- Davidson EA, Chorover J, Dail DB. A mechanism of abiotic immobilization of nitrate in forest ecosystems: The ferrous wheel hypothesis. *Global Change Biology*. 2003;9:228–236. [Google Scholar]
- De Wit HA, Hindar A, Hole L. Winter climate affects long-term trends in stream water nitrate in acid-sensitive catchments in southern Norway. *Hydrology and Earth System Sciences*. 2008; 12:393–403. [Google Scholar]
- Deegan LA, Johnson DS, Warren RS, Peterson BJ, Fleeger JW, Fagherazzi S, Wollheim WM. Coastal eutrophication as a driver of salt marsh loss. *Nature*. 2012;490:388–392. [PubMed] [Google Scholar]
- Dias T, Malveiro S, Martins-Loução MA, Sheppard LJ, Cruz C. Linking increased N-driven biodiversity changes with soil N availability in a Mediterranean-Type Ecosystem. *Plant and Soil*. 2011;341:125–136. [Google Scholar]
- Dise N, Wright R. Nitrogen leaching from European forests in relation to nitrogen deposition. *Forest Ecology and Management*. 1995;71:153–161. [Google Scholar]
- Driscoll CT, Whitall D, Aber J, Boyer E, Castro M, Cronan C, Goodale CL, Groffman P, et al. Nitrogen Pollution in the Northeastern United States: Sources, Effects, and Management Options. *BioScience*. 2003;53:357–374. [Google Scholar]
- Driscoll CT, Lambert KF, Chapin FS, III, Nowak DJ, Spies TA, Swanson FJ, Kittredge DB, Hart CM. Science and society: The role of long-term studies in environmental stewardship. *BioScience*. 2012;62:354–366. [Google Scholar]
- Eimers MC, Buttle JM, Watmough SA. The contribution of rain-on-snow events to annual $\text{NO}_3\text{-N}$ export from a forested catchment in south-central Ontario, Canada. *Applied Geochemistry*. 2007;22:1105–1110. [Google Scholar]
- Evans CD, Cooper DM, Monteith DT, Helliwell RC, Moldan F, Hall J, Rowe EC, Cosby BJ. Linking monitoring and modelling: Can long-term datasets be used more effectively as a basis for large-scale prediction? *Biogeochemistry*. 2010;101:211–227. [Google Scholar]
- Fang Y, Gundersen P, Vogt RD, Koba K, Chen F, Chen XY, Yoh M. Atmospheric deposition and leaching of nitrogen in Chinese forest ecosystems. *Journal of Forest Research*. 2011;16:341–350. [Google Scholar]
- Fernandez IJ, Adams MB, SanClements MD, Norton SA. Comparing decadal responses of whole-watershed manipulations at the Bear Brook and Fernow experiments. *Environmental Monitoring and Assessment*. 2010;171:149–161. [PubMed] [Google Scholar]

Fowler D, Smith R, Muller J, Cape JN, Sutton M, Erisman JW, Fagerli H. Long term trends in sulphur and nitrogen deposition in Europe and the cause of non-linearities. *Water, Air, & Soil Pollution: Focus*. 2007;7:41–47. [Google Scholar]

Freppaz M, Ceili L, Marcheilli M, Zanini E. Snow removal and its influence on temperature and N dynamics in alpine soils (Vallée d'Aoste, northwest Italy) *Journal of Plant Nutrition and Soil Science*. 2008;171:672–680. [Google Scholar]

Galloway JN, Aber JD, Erisman JW, Seitzinger SP, Howarth RW, Cowling EB, Cosby BJ. The nitrogen cascade. *BioScience*. 2003;53:341–356. [Google Scholar]

Galloway JN, Dentener FJ, Capone DG, Boyer EW, Howarth RW, Seitzinger SP, Asner GP, Cleveland CC, et al. Nitrogen Cycles: Past, Present and Future. *Biogeochemistry*. 2004;70:153–226. [Google Scholar]

Galloway JN, Townsend AR, Erisman JW, Bekunda M, Cai Z, Freney JR, Martinelli LA, Seitzinger SP, et al. Transformation of the Nitrogen Cycle: Recent Trends, Questions, and Potential Solutions. *Science*. 2008;320:889–892. [PubMed] [Google Scholar]

Gärdenäs AI, Ågren GI, Bird JA, Clarholm M, Hallin S, Ineson P, Kätterer T, Knicker H, Nilsson SI, et al. Knowledge gaps in soil carbon and nitrogen interactions – From molecular to global scale. *Soil Biology & Biochemistry*. 2011;43:702–717. [Google Scholar]

George DG, Maberly SC, Hewitt DP. The influence of the North Atlantic Oscillation on the physical, chemical and biological characteristics of four lakes in the English Lake District. *Freshwater Biology*. 2004;49:760–774. [Google Scholar]

Groffman P, Butterbach-Bahl K, Fulweiler RW, Gold AJ, Morse JL, Stander EK, Tague C, Tonitto C, et al. Challenges to incorporating spatially and temporally explicit phenomena (hotspots and hot moments) in denitrification models. *Biogeochemistry*. 2009;92:49–77. [Google Scholar]

Groffman PM, Rustad LE, Templer PH, Campbell JL, Christenson LM, Lany NK, Socci AM, Vadeboncoeur MA, et al. Long-term integrated studies show complex and surprising effects of climate change in the northern hardwood forest. *BioScience*. 2012;62:1056–1066. [Google Scholar]

Gruber N, Galloway JN. An Earth-system perspective of the global nitrogen cycle. *Nature*. 2008;451:293–296. [PubMed] [Google Scholar]

Gundersen P, Emmett BA, Kjønaas OJ, Koopmans CJ, Tietemad A. Impact of nitrogen deposition on nitrogen cycling in forests: A synthesis of NITREX data. *Forest Ecology and Management*. 1998;101:37–55. [Google Scholar]

Gupta SK, Gupta RC, Seth AK. Methaemoglobinaemia in areas with high nitrate concentration in drinking water. *National Medical Journal of India*. 2000;13:58–61. [PubMed] [Google Scholar]

Harris TD, Wilhelm FM, Graham JL, Loftin KA. Experimental manipulation of TN:TP ratios suppress cyanobacterial biovolume and microcystin concentration in large-scale in situ microcosms. *Lake Reserv Manag.* 2014;30:72–83. [Google Scholar]

Hedwall PO, Nordin A, Strengbom J, Brunet J, Olsson B. Does background nitrogen deposition affect the response of boreal vegetation to fertilization? *Oecologia.* 2013;173:615–624. [PubMed] [Google Scholar]

Hoben, J.P., R.J. Gehl, N. Millar, P.R. Grace, and G.P. Robertson. 2011. Nonlinear nitrous oxide (N₂O) response to nitrogen fertilizer in on-farm corn crops of the US Midwest. *Global Change Biology* 17: 1140–1152.

Högberg MN, Chen Y, Högberg P. Gross nitrogen mineralisation and fungi-to-bacteria ratios are negatively correlated in boreal forests. *Biology and Fertility of Soils.* 2007;44:363–366. [Google Scholar]

Högberg MN, Briones MJI, Keel SG, Metcalfe DB, Campbell C, Midwood AJ, Thornton B, Hurry V, et al. Quantification of effects of season and nitrogen supply on tree below-ground carbon transfer to ectomycorrhizal fungi and other soil organisms in a boreal pine forest. *New Phytologist.* 2010;187:485–493. [PubMed] [Google Scholar]

Howarth RW, Marino R. Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: Evolving views over three decades. *Limnology and Oceanography.* 2006;51:364–376. [Google Scholar]

Howarth R, Swaney D, Billen G, Garnier J, Hong B, Humborg C, Johnes P, Mörtz C-M, et al. Nitrogen fluxes from the landscape are controlled by net anthropogenic nitrogen inputs and by climate. *Frontiers in Ecology and the Environment.* 2011;10:37–43. [Google Scholar]

IPCC. 2007. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge: Cambridge University Press.

Jetten MS, Cirpus I, Kartal B, van Niftrik L, van de Pas-Schoonen KT, Sliekers O, Haaijer S, van der Star W, Schmid M, et al. 1994-2004: 10 years of research on the anaerobic oxidation of ammonium. *Biochemical Society Transactions.* 2005;33:119–123. [PubMed] [Google Scholar]

Johansson O, Palmqvist K, Olofsson J. Nitrogen deposition drives lichen community changes through differential species responses. *Glob Chang Biol.* 2012;18:2626–2635. [Google Scholar]

Jovan S, Riddell J, Padgett PE, Nash TH., III Eutrophic lichens respond to multiple forms of N: Implications for critical levels and critical loads research. *Ecological Applications.* 2012;22:1910–1922. [PubMed] [Google Scholar]

Kaste Ø, Austnes K, Vestgarden LS, Wright RF. Manipulation of snow in small headwater catchments at Storgama, Norway: Effects on leaching of inorganic nitrogen. *AMBIO.* 2008;37:29–37. [PubMed] [Google Scholar]

Knorr M, Frey SD, Curtis PS. Nitrogen additions and litter decomposition: A meta-analysis. *Ecology*. 2005;86:3252–3257. [Google Scholar]

Kopáček, J., and M. Posch. 2011. Anthropogenic nitrogen emissions during the Holocene and their possible effects on remote ecosystems. *Global Biogeochemical Cycles* 25: GB2017. doi:10.1029/2010GB003779.

Kopáček J, Hejzlar J, Posch M. Factors controlling the export of nitrogen from agricultural land in a large central European catchment during 1900–2010. *Environmental Science and Technology*. 2013;47:6400–6407. [PubMed] [Google Scholar]

Kopáček J, Hejzlar J, Posch M. Quantifying nitrogen leaching from diffuse agricultural and forest sources in a large heterogeneous catchment. *Biogeochemistry*. 2013;115:149–165. [PubMed] [Google Scholar]

Kopáček J, Cosby BJ, Evans CD, Hruška J, Moldan F, Oulehle F, Šantrůčková H, Tahovská K, et al. Nitrogen, organic carbon and sulphur cycling in terrestrial ecosystems: Linking nitrogen saturation to carbon limitation of soil microbial processes. *Biogeochemistry*. 2013;115:33–51. [Google Scholar]

Kurian LM, Lautz LK, Mitchell MJ. Winter hydrology and NO₃[–] concentrations in a forested watershed: A detailed field study in the Adirondack Mountains of New York. *Journal of the American Water Resources Association*. 2012;49:264–283. [Google Scholar]

Landberg JO, Carlstrom M, Larsen FJ, Weitzberg E. Roles of dietary inorganic nitrate in cardiovascular health and disease. *Cardiovascular Research*. 2011;89:525–532. [PubMed] [Google Scholar]

Lassaletta L, Billen G, Grizzetti B, Garnier J, Leach AM, Galloway JN. Food and feed trade as a driver in the global nitrogen cycle: 50-year trends. *Biogeochemistry*. 2014;118:225–241. [Google Scholar]

Leach AM, Galloway JN, Bleeker A, Erisman JW, Kohn R, Kitzes J. A nitrogen footprint model to help consumers understand their role in nitrogen losses to the environment. *Environmental Development*. 2012;1:40–66. [Google Scholar]

Likens GE, Driscoll CT, Buso DC. Long-term effects of acid rain: Response and recovery of a forest ecosystem. *Science*. 1996;272:244–246. [Google Scholar]

Löfgren S, Aastrup M, Bringmark L, Hultberg H, Lewin-Pihlblad L, Lundin L, Karlsson GP, Thunholm B. Recovery of Soil Water, Groundwater, and Streamwater from Acidification at the Swedish Integrated Monitoring Catchments. *AMBIO*. 2011;40:836–856. [PMC free article] [PubMed] [Google Scholar]

Maberly SC, King L, Dent MM, Jones RI, Gibson CE. Nutrient limitation of phytoplankton and periphyton growth in upland lakes. *Freshwater Biology*. 2002;47:2136–2152. [Google Scholar]

- Makoto, K., T. Kajimoto, L. Koyama, G. Kudo, H. Shibata, Y. Yanai, and J.H.C. Cornelissen. 2013 Winter climate change in plant–soil systems: Summary of recent findings and future perspectives. *Ecological Research* (in press).
- Matson PA, McDowell WH, Townsend AR, Vitousek PM. The globalization of N deposition: Ecosystem consequences in tropical environments. *Biogeochemistry*. 1999;46:67–83. [Google Scholar]
- Millennium Ecosystem Assessment Board. 2005. *Ecosystems and human well-being: Current state and trends*, Vol. 1. Washington D.C.: Island Press.
- Mitchell MJ. Nitrate dynamics of forested watersheds: Spatial and temporal patterns in North America, Europe and Japan. *Journal of Forest Research*. 2011;16:333–340. [Google Scholar]
- Mitchell MJ, Driscoll CT, Kahl JS, Likens GE, Murdoch PS, Pardo LH. Climatic control of nitrate loss from forested watersheds in the northeast United States. *Environmental Science and Technology*. 1996;30:2609–2612. [Google Scholar]
- Monson RK, Lipson DL, Burns SP, Turnipseed AA, Delany AC, Williams MW, Schmidt SK. Winter forest soil respiration controlled by climate and microbial community composition. *Nature*. 2006;439:711–714. [PubMed] [Google Scholar]
- Monteith DT, Evans CD, Reynolds B. Are temporal variations in the nitrate content of UK upland freshwaters linked to the North Atlantic Oscillation? *Hydrological Processes*. 2000;14:1745–1749. [Google Scholar]
- Monteith DT, Evans CD, Henrys PA, Simpson GL, Malcolm IA. Trends in the hydrochemistry of acid-sensitive surface waters in the UK 1988–2008. *Ecological Indicators*. 2014;37:287–303. [Google Scholar]
- Niu S, Wu M, Han Y, Xia J, Zhang Z, Yang H, Wan S. Nitrogen effects on net ecosystem carbon exchange in a temperate steppe. *Global Chang Biol*. 2010;16:144–155. [Google Scholar]
- Noe GB, Hupp CR. Carbon, nitrogen, and phosphorus accumulation in floodplains of Atlantic coastal plain rivers, USA. *Ecological Applications*. 2005;15:1178–1190. [Google Scholar]
- Ochoa-Hueso R, Allen EB, Branquinho C, Cruz C, Dias T, Fenn ME, Manrique E, Pérez-Corona ME, et al. Nitrogen deposition effects on Mediterranean-type ecosystems: An ecological assessment. Review. *Environmental Pollution*. 2011;159:2265–2279. [PubMed] [Google Scholar]
- Ohte N, Mitchell MJ, Shibata H, Tokuchi N, Toda H, Iwatsubo G. Comparative evaluation on nitrogen saturation of forest catchments in Japan and northeastern United States. *Water, Air, and Soil pollution*. 2001;131:649–654. [Google Scholar]
- Oulehle F, McDowell WH, Aitkenhead-Peterson JA, Krám P, Hruška J, Navrátil T, Buzek F, Fottová D. Long-Term Trends in Stream Nitrate Concentrations and Losses Across

Watersheds Undergoing Recovery from Acidification in the Czech Republic. *Ecosystems*. 2008;11:410–425. [Google Scholar]

Pardo LH, Fenn ME, Goodale CL, Geiser LH, Driscoll CT, Allen EB, Baron JS, Bobbink R, et al. Effects of nitrogen deposition and empirical nitrogen critical loads for ecoregions of the United States. *Ecological Applications*. 2011; 21:3049–3082. [Google Scholar]

Park J, Mitchell MJ, McHale PJ, Christopher SF, Myers TP. Interactive effects of changing climate and atmospheric deposition on N and S biogeochemistry in a forested watershed of the Adirondack Mountains, New York State. *Global Change Biology*. 2003;9:1602–1619. [Google Scholar]

Peñuelas J, Prieto P, Beier C, Cesaraccio C, De Angelis P, De Dato G, Emmett BA, Estiarte M, et al. Response of plant species richness and primary productivity in shrublands along a north–south gradient in Europe to seven years of experimental warming and drought: Reductions in primary productivity in the heat and drought year of 2003. *Glob Chang Biol*. 2007;13:2563–2581. [Google Scholar]

Pereira HM, Ferrier S, Walters M, Geller GN, Jongman RHG, Scholes RJ, Bruford MW, Brummitt N, et al. Essential Biodiversity Variables. *Science*. 2013;339:277–278. [PubMed] [Google Scholar]

Piatek KB, Mitchell MJ, Silva SR, Kendall C. Sources of nitrate in snowmelt discharge: Evidence from water chemistry and stable isotopes of nitrate. *Water, Air, and Soil pollution*. 2005;165:13–35. [Google Scholar]

Pinho P, Dias T, Cruz C, Tang YS, Sutton MA, Martins-Loução M-A, Máguas C, Branquinho C. Using lichen functional-diversity to assess the effects of atmospheric ammonia in Mediterranean woodlands. *Journal of Applied Ecology*. 2011;48:1107–1116. [Google Scholar]

Pinho P, Theobald MR, Dias T, Tang YS, Cruz C, Martins-Loucao MA, Maguas C, Sutton M, et al. Critical loads of nitrogen deposition and critical levels of atmospheric ammonia for semi-natural Mediterranean evergreen woodlands. *Biogeosciences*. 2012;9:1205–1215. [Google Scholar]

Pregitzer KS, Burton AJ, Zak DR, Talhelm AF. Simulated chronic nitrogen deposition increases carbon storage in Northern Temperate forests. *Global Change Biology*. 2008;14:142–153. [Google Scholar]

Rask M, Arvola L, Forsius M, Vuorenmaa J. Preface to the special issue “Integrated monitoring in the Valkea-Kotinen catchment during 1990–2009: Abiotic and biotic responses to changes in air pollution and climate. *Boreal Environment Research*. 2014;19:1–3. [Google Scholar]

Reay DS, Davidson EA, Smith KA, Smith P, Melillo JM, Dentener F, Crutzen PJ. Global agriculture and nitrous oxide emissions. *Nature Climate Change*. 2012;2:410–416. [Google Scholar]

Rissanen A, Tirola M, Ojala A. Spatial and temporal variation in denitrification and in the denitrifier community in a boreal lake. *Aquatic Microbial Ecology*. 2011;64:27–40. [Google Scholar]

Robertson GP, Paul EA, Harwood RR. Greenhouse Gases in Intensive Agriculture: Contributions of Individual Gases to the Radiative Forcing of the Atmosphere. *Science*. 2000;289:1922–1925. [PubMed] [Google Scholar]

Robertson GP, Collins SL, Foster DR, Brokaw N, Ducklow HW, Gragson TL, Gries C, Hamilton SK, et al. Long-Term Ecological Research in a Human-Dominated World. *BioScience*. 2012;62:342–353. [Google Scholar]

Rogora M, Arese C, Balestrini R, Marchetto A. Climate control on sulphate and nitrate concentrations in alpine streams of Northern Italy along a nitrogen saturation gradient. *Hydrology and Earth System Sciences*. 2008;12:371–381. [Google Scholar]

Sanchez P, Palm C, Sachs J, Denning G, Flor R, Harawa R, Jama B, Kiflemariam T, et al. The African millennium villages. *Proceedings of the National Academy of Sciences of the United States of America*. 2007;104:16775–16780. [PMC free article] [PubMed] [Google Scholar]

Schubert H, Wasmund N. Das Phytoplankton des Strelasundes und des Kubitzer Boddens. *Meer Museum*. 2005;18:83–92. [Google Scholar]

Schubert, H., N. Wasmund, and K. Sellner. 2010. Long-term investigations in brackish ecosystems. In *Long-term ecological research*, ed. F. Müller, C. Baessler, H. Schubert, and S. Klotz, 163–178: Dordrecht: Springer.

Shaver GR, Johnson LC, Cades DH, Murray G, Laundre JA, Rastetter EB, Nadelhoffer KJ, Giblin AE. Biomass and CO₂ flux in wet sedge tundras: Responses to nutrients, temperature, and light. *Ecological Monographs*. 1998;68:75–97. [Google Scholar]

Shibata H, Sugawara O, Toyoshima H, Wondzell SM, Nakamura F, Kasahara T, Swanson FJ, Sasa K. Nitrogen dynamics in the hyporheic zone of a forested stream during a small storm, Hokkaido, Japan. *Biogeochemistry*. 2004;69:83–104. [Google Scholar]

Shibata H, Hasegawa Y, Watanabe T, Fukuzawa K. Impact of snowpack decrease on net nitrogen mineralization and nitrification in forest soil of northern Japan. *Biogeochemistry*. 2013;116:69–82. [Google Scholar]

Skiba A, Smith KA. The control of nitrous oxide emissions from agricultural and natural soils. *Chemosphere—Global Change Science*. 2000;2:379–386. [Google Scholar]

Spannhake EW, Reddy SPM, Jacoby DB, Yu XY, Saatian B, Tian J. Synergism between rhinovirus infection and oxidant pollutant exposure enhances airway epithelial cell cytokine production. *Environmental Health Perspectives*. 2002; 110:665–670. [PMC free article] [PubMed] [Google Scholar].

Stal LJ, Zehr JP. Cyanobacterial nitrogen fixation in the ocean: Diversity, regulation and ecology. In: Herrero A, Flores E, editors. *The cyanobacteria: Molecular biology, genomics and evolution*. Caister: Academic Press; 2008. pp. 423–446. [Google Scholar].

Stoddard JL. Long-term changes in watershed retention of nitrogen: Its causes and aquatic consequences. In: Baker LA, editor. *Environmental chemistry of lakes and reservoirs*. Washington, DC: American Chemical Society; 1994. pp. 223–284. [Google Scholar].

Suddick, E.C., and E.A. Davidson. 2012. *The Role of Nitrogen in Climate Change and the Impacts of Nitrogen-Climate Interactions on Terrestrial and Aquatic Ecosystems, Agriculture, and Human Health in the United States: A Technical Report Submitted to the US National Climate Assessment*. North American Nitrogen Center of the International Nitrogen Initiative (NANC-INI), Woods Hole Research Center, 149 Woods Hole Road, Falmouth, MA.

Sutton MA, Howard CM, Erisman JW, Billen G, Bleeker A, Grennfelt P, van Grinsven H, Grizzetti B. *The European Nitrogen Assessment*. Cambridge: Cambridge University Press; 2011. [Google Scholar].

Thornton PE, Doney SC, Lindsay K, Moore JK, Mahowald N, Randerson JT, Fung I, Lamarque J-F, et al. Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: Results from an atmosphere-ocean general circulation model. *Biogeosciences*. 2009;6:2099–2120. [Google Scholar].

Tierney GT, Fahey TJ, Groffman PM, Hardy JP, Fitzhugh RD, Driscoll CT. Soil freezing alters fine root dynamics in a northern hardwood forest. *Biogeochemistry*. 2001;56:175–190. [Google Scholar]

Urakawa R, Toda H, Haibara K, Aiba Y. Long-term hydrochemical monitoring in an Oyasan Experimental Forest Watershed comprised of two small forested watersheds of Japanese cedar and Japanese cypress. *Ecol Res (Data paper)* 2012;27:245. [Google Scholar]

Van Groenigen JW, Velthof GL, Oenema O, Van Groenigen KJ, Van Kessel C. Towards an agronomic assessment of N₂O emissions: A case study for arable crops. *European Journal of Soil Science*. 2010;61:903–913. [Google Scholar]

Van Maanen JMS, Welle IJ, Hageman G, Dallinga JW, Mertens PL, Kleinjans JC. Nitrate contamination of drinking water: Relationship with HPRT variant frequency in lymphocyte DNA and urinary excretion of N-nitrosamines. *Environmental Health Perspectives*. 1996;104:522–528. [PMC free article] [PubMed] [Google Scholar]

Vermaat, J.E., S. Broekx, B. Van Eck, G. Engelen, F. Hellmann, J.L. De Kok, H. Van der Kwast, J. Maes, W. Salomons, et al. 2012. Nitrogen source apportionment for the catchment, estuary, and adjacent coastal waters of the river Scheldt. *Ecology and Society* 17: 30. doi:10.5751/ES-04889-170230.

Vitousek, P.M., R. Naylor, T. Crews, M.B. David, L.E. Drinkwater, E. Holland, P.J. Johnes, and J. Katzenberger. 2009. Nutrient imbalances in agricultural development. *Science* 324: 1519–1520. [PubMed]

Von Mutius E. Current review of allergy and immunology. *Journal of Allergy and Clinical Immunology*. 2000; 105:9–19. [PubMed] [Google Scholar]

Wallenstein MD, Myrold DD, Firestone M, Voytek M. Environmental controls on denitrifying communities and denitrification rates: Insights from molecular methods. *Ecological Applications*. 2006; 16:2143–2152. [PubMed] [Google Scholar]

Ward MH, Mark SD, Cantor KP, Weisenburger DD, Correa-Villaseñor A, Zahm SH. Drinking water nitrate and the risk of non-Hodgkin's lymphoma. *Epidemiology*. 1996;7:465–471. [PubMed] [Google Scholar]

Weyer PJ, Cerhan J, Kross BC, Hallberg GR, Kantamneni J, Breuer G, Jones MP, Zheng W, et al. Municipal drinking water nitrate level and cancer risk in older women: The Iowa Women's Health Study. *Epidemiology*. 2001;12:327–338. [PubMed] [Google Scholar]

Wipf S, Rixen C. A review of snow manipulation experiments in Arctic and alpine tundra ecosystems. *Polar Research*. 2010;29:95–109. [Google Scholar]

Wright RF. Effect of increased carbon dioxide and temperature on runoff chemistry at a forested catchment in southern Norway (CLIMEX Project) *Ecosystems*. 1998;1:216–225. [Google Scholar]

Zaehle S, Ciais P, Friend AD, Prieur V. Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions. *Nature Geoscience*. 2011;4:601–605. [Google Scholar]